

**Centre for AIDS Reagents**  
**Data Sheet**

<b>Reagent:</b>	Anti-SIVmac251 Nef Monoclonal (17.2)
<b>Catalogue:</b>	<b>100400</b>
<b>Lot Number:</b>	150215
<b>Provided:</b>	250 µg affinity purified antibody at 1 mg/mL in PBS pH 7.4, with 0.1% sodium azide.
<b>Description:</b>	This mAb targets SIVmac251 Nef. Mapped to amino acids 71-80 (-PAEEREKLAY-, SIV).
<b>Host:</b>	Balb/c splenocyte x SP2 myeloma.
<b>Titer:</b>	ELISA titer: 0.05 to 0.1 µg/ml , WB: 1 µg/ml. End users should determine optimal concentrations to be used in assays.
<b>Special characteristics:</b>	Mice were immunized with MS2 phage RNA polymerase conjugated to SIVmac251 Nef. Reacts in ELISA, WB, RIP, IHC with Nef of SIVmac251 strain. Reacts in IF with Nef of SIVmac239 strain and SIVmac1A11 strain.  Purified from E. coli using Protein G. Alternate names: Clone 17
<b>Recommended Storage:</b>	Keep at 4°C for short term storage and -80°C for long term storage. Avoid freeze-thaw cycles as reagent degradation may result.
<b>Contributor:</b>	Dr. Kai Krohn and Dr. Vladimir Ovod.
<b>Isotype:</b>	IgG <sub>1</sub>
<b>References:</b>	Piguet V, Chen Y-L, Mangasarian A, Foti M, Carpentier J-L, Trono D. (1998) Mechanism of Nef-induced CD4 endocytosis: Nef connects CD4 with the µ chain of adaptor complexes. EMBO, 17 (9): 2472-81.  Chan KS, Verardi PH, Legrand FA, Yilma TD. (2005) Nef from pathogenic simian immunodeficiency virus is a negative factor for vaccinia virus. PNAS 102(24): 8734-9.

**ACKNOWLEDGEMENTS:**

Publications should acknowledge the contributor and the Centre for AIDS Reagents. Acknowledgments should read: "The *Name of Reagent (Repository Number)* was obtained from the Centre for AIDS Reagents, NIBSC, UK, supported by EURIPRED (EC FP7 INFRASTRUCTURES-2012 - INFRA-2012-1.1.5.: Grant Number 31266). [www.euripred.eu/](http://www.euripred.eu/)