

Centre For AIDS Reagents

Data Sheet

REAGENT:	J-Lat Tat-GFP Cells (H2)
REPOSITORY REFERENCE:	100 947
PROVIDED:	1 mL at 7.7×10^6 cells. Viability = 79%.
LOT NUMBER:	150246
RELEASE RESTRICTION:	NIH category C
STORAGE:	Liquid nitrogen
DESCRIPTION:	These cells are Jurkat cells that bear the integrated retroviral construct LTR-Tat-IRES-GFP.
SPECIAL CHARACTERISTICS:	Jurkat cells were infected with viral particles bearing the retroviral construct LTR-Tat-IRES-GFP. Cells that were GFP negative, but could be stimulated to express GFP were selected. For the other similar cell, please see 100941-100948.
CELL TYPE:	Jurkat - T lymphocyte cell line
FREEZE MEDIUM:	FBS, 90%; DMSO, 10%.
GROWTH CHARACTERISTIC:	No special requirements, split 1:3 at 1×10^6 cells/ml. Cells grow in suspension, usually singly but some clumping has been noted.
PROPAGATION MEDIUM:	RPMI 1640, 90%; FBS, 10%; supplemented with penicillin G (100 U/ml), streptomycin (100 µg/ml), L-glutamine (2 mM, 0.3 mg/ml).
MORPHOLOGY:	Small, spherical cells in suspension. Morphology usually does not vary.
CONTRIBUTOR:	Dr. Eric Verdin.
REFERENCES:	Jordan A, Bisgrove D, Verdin E. HIV reproducibly establishes a latent infection after acute infection of T cells in vitro. <i>EMBO J</i> 22 :1868-1877, 2003. Jordan A, Defechereux P, Verdin E. The site of HIV-1 integration in the human genome determines basal transcriptional activity and response to Tat transactivation. <i>EMBO J</i> 20 :1726-1738, 2001.

ACKNOWLEDGMENT:

The following reagent was obtained by CFAR – NIBSC via the NIH AIDS Reagent Program, Division of AIDS, NIAID, NIH: J-Lat Tat-GFP (clone #) from Dr. Eric Verdin. Also include the references cited above in any publication.

Please ensure that you send us a copy of any papers Resulting from work using reagents acquired through CFAR, this can be by e-mail or printed copy.